

Appendix B: Outcome Measures

Domain	Name	Screening (S) or Assessment (A)	Time to Complete
Measures of Stroke Severity (Acute care only)	Orpington Prognostic Scale	S & A	5 min
	National Institute of Health Stroke Scale (NIHSS)	A	< 10 min
	Canadian Neurological Scale (CNS)	A	5 min
	Glasgow Coma Scale	A	< 5 min
Medical co-morbidities:	Charlson Co-morbidity Index	S	5 min
Cognition	Montreal Cognitive Assessment (MoCA[®]) (cross continuum)	S	10-15
Visual Perception	Line Bisection Test	S	< 5 min
	Motor Free Visual Perceptual Test (MVPT)	A	15-20 min
Self Care Activities of Daily Living	AlphaFIM[®] Instrument (acute only)	A	15-20
	FIM[®] Instrument (Rehab & Community)	A	30-45
Mobility & Function	Berg Balance Scale (cross continuum)	A	10-15 min
	Timed Up and Go (TUG)	S	1-2 min
	Chedoke-McMaster Stroke Assessment (CMSA) (cross continuum)	S & A	45-60 min
Depression	Hospital Anxiety Depression Scale (HADS)	S	2-5 min
	Montgomery Asberg Depression Rating Scale (MADRS)	A	20-60
Language	Boston Diagnostic Aphasia Examination (BDAE)	A	40-60 min (short version)
	Frenchay Aphasia Screening Test (FAST)	S	5 min
Swallowing	Toronto Bedside Stroke Screening Tool (ToRBSST [®])	S & A	10 min
Participation	Stroke Impact Scale (SIS)	A	15-20 min
	Reintegration to Normal Living Index (RNLI)	A	10 min
Pain	Visual Analogue Scale	A	5 min
	Pain numerical scale	A	5 min

Those tools in red represent core screening and assessment tools recommended by the Toronto Stroke Networks